

On the relation between the full Kostant–Toda lattice and multiple orthogonal polynomials

D. Barrios Rolanía^a, A. Branquinho^{b,*}, A. Foulquié Moreno^c

^a Facultad de Informática, Universidad Politécnica de Madrid, 28660 Boadilla del Monte, Spain

^b CMUC, Department of Mathematics, University of Coimbra, Largo D. Dinis, 3001-454 Coimbra, Portugal

^c Universidade de Aveiro, Campus de Santiago 3810, Aveiro, Portugal

A B S T R A C T

The correspondence between a high-order non-symmetric difference operator with complex coefficients and the evolution of an operator defined by a Lax pair is established. The solution of the discrete dynamical system is studied, giving explicit expressions for the resolvent function and, under some conditions, the representation of the vector of functionals, associated with the solution for our integrable systems. The method of investigation is based on the evolutions of the matricial moments.

1. Introduction

We consider the following special *full Kostant–Toda* system,

$$\left. \begin{aligned} \dot{a}_n &= b_n - b_{n-1}, \\ \dot{b}_n &= b_n(a_{n+1} - a_n) + c_n - c_{n-1}, \\ \dot{c}_n &= c_n(a_{n+2} - a_n), \end{aligned} \right\} \quad n \in \mathbb{N}, \quad (1)$$

where the dot means differentiation with respect to $t \in \mathbb{R}$ and we assume $b_0 \equiv 0$, $c_0 \equiv 0$, and $c_n \neq 0$. It is well known that these equations can be written as a *Lax pair* $\dot{J} = [J, J_-]$, where $[M, N] = MN - NM$ is the *commutator* of the operators M and N , and J, J_- are the operators which matrix representation is given, respectively, by

$$J = \begin{pmatrix} a_1 & 1 & & \\ b_1 & a_2 & 1 & \\ c_1 & b_2 & a_3 & \ddots \\ 0 & c_2 & b_3 & \ddots \\ & \ddots & \ddots & \ddots \end{pmatrix}, \quad J_- = \begin{pmatrix} 0 & & & \\ b_1 & 0 & & \\ c_1 & b_2 & 0 & \\ 0 & c_2 & b_3 & \ddots \\ & \ddots & \ddots & \ddots \end{pmatrix}. \quad (2)$$

Here, and in the following, we suppress the explicit t -dependence for brevity. Also, we identify an operator and its matrix representation with respect to the canonical basis. For the sake of simplicity, we only consider a four banded matrix J in this work, but the method can be extended to higher order banded matrices J .

* Corresponding author.

E-mail addresses: dbarrios@fi.upm.es (D. Barrios Rolanía), ajplb@mat.uc.pt (A. Branquinho), foulquie@ua.pt (A. Foulquié Moreno).

In [1,8] the authors considered some special cases of the systems studied here, and in [4] some finite full Kostant–Toda systems are considered and solved using bi-orthogonal systems of polynomials.

When J is a bounded operator, then it is possible to define the *resolvent operator*,

$$(zI - J)^{-1} = \sum_{n \geq 0} \frac{J^n}{z^{n+1}}, \quad |z| > \|J\| \quad (3)$$

(see [12, Theorem 3, p. 211]). We denote by M_{ij} the 2×2 block, of any infinite matrix M , formed by the entries of rows $2i - 1, 2i$ and columns $2j - 1, 2j$. In this way, for each $n \in \mathbb{N}$, J^n can be written as a blocked matrix,

$$J^n = \begin{pmatrix} J_{11}^n & J_{12}^n & \cdots \\ J_{21}^n & J_{22}^n & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}. \quad (4)$$

Moreover, we define $\mathcal{R}_J(z)$ as the main block of $(zI - J)^{-1}$, this is, $\mathcal{R}_J(z) := (zI - J)_{11}^{-1}$. Then, from (3) we have

$$\mathcal{R}_J(z) = \sum_{n \geq 0} \frac{J_{11}^n}{z^{n+1}}, \quad |z| > \|J\|. \quad (5)$$

As a consequence of the Lax pair representation, for (1) and other systems, the operator theory is a useful tool and a remarkable connection between the integrable systems and the approximation theory. Consider the sequence of polynomials $\{P_n\}$ given by the recurrence relation

$$\left. \begin{aligned} c_{n-1}P_{n-2}(z) + b_nP_{n-1}(z) + (a_{n+1} - z)P_n(z) + P_{n+1}(z) &= 0, \quad n = 0, 1, \dots, \\ P_0(z) &= 1, \quad P_{-1}(z) = P_{-2}(z) = 0. \end{aligned} \right\} \quad (6)$$

Taking $\mathcal{B}_n(z) = (P_{2n}(z), P_{2n+1}(z))^T$, we can rewrite (6) as

$$\left. \begin{aligned} C_n \mathcal{B}_{n-1}(z) + (B_{n+1} - zI_2) \mathcal{B}_n(z) + A \mathcal{B}_{n+1}(z) &= 0, \quad n = 0, 1, \dots, \\ \mathcal{B}_{-1}(z) &= 0, \quad \mathcal{B}_0(z) = (1, z - a_1)^T \end{aligned} \right\} \quad (7)$$

where

$$A = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad C_n = \begin{pmatrix} c_{2n-1} & b_{2n} \\ 0 & c_{2n} \end{pmatrix}, \quad B_n = \begin{pmatrix} a_{2n-1} & 1 \\ b_{2n-1} & a_{2n} \end{pmatrix}, \quad n \in \mathbb{N}, \quad (8)$$

and C_0 is an arbitrary (2×2) -matrix. In (7) and in the following, we denote for the sake of simplicity $(0, 0)^T$ and $0 \in \mathbb{R}$ in the same way. We recall that the polynomials $P_n(z)$ and the vectorial polynomials $\mathcal{B}_n(z)$ depend on $t \in \mathbb{R}$, when this dependence holds for the coefficients a_n, b_n, c_n .

Our main goal is to study the solutions of (1) in terms of the operator J and its associated vectorial polynomials $\mathcal{B}_n(z)$. We underline that, in a different context (cf. [2, Theorem 2]), the characterization of solutions of an integrable system was established in terms of the derivative of the polynomials associated with J . In this work, using the sequence of vectorial polynomials $\{\mathcal{B}_n\}$, we extend that result. Our first result is the following.

Theorem 1. Assume that the sequence $\{a_n, b_n, c_n\}$, $n \in \mathbb{N}$, is uniformly bounded, i.e. there exists $K \in \mathbb{R}_+$ such that $\max\{|a_n(t)|, |b_n(t)|, |c_n(t)|\} \leq M$ for all $n \in \mathbb{N}$ and $t \in \mathbb{R}$. Assume, also, $c_n(t) \neq 0$ for all $n \in \mathbb{N}$ and $t \in \mathbb{R}$. Then, the following conditions are equivalent:

(a) $\{a_n, b_n, c_n\}$, $n \in \mathbb{N}$, is a solution of (1), this is,

$$\dot{J} = [J, J_-]. \quad (9)$$

(b) For each $n \in \mathbb{N} \cup \{0\}$ we have

$$\frac{d}{dt} J_{11}^n = J_{11}^{n+1} - J_{11}^n B_1 + [J_{11}^n, (J_-)_{11}]. \quad (10)$$

(c) For all $z \in \mathbb{C}$ such that $|z| > \|J\|$,

$$\dot{\mathcal{R}}_J(z) = \mathcal{R}_J(z)(zI_2 - B_1) - I_2 + [\mathcal{R}_J(z), (J_-)_{11}], \quad (11)$$

where $\mathcal{R}_J(z)$ is given by (5).

(d) For each $n \in \mathbb{N} \cup \{0\}$, the polynomial \mathcal{B}_n defined by (7) satisfies

$$\dot{\mathcal{B}}_n(z) = -C_n \mathcal{B}_{n-1}(z) - D_n \mathcal{B}_n(z), \quad (12)$$

$$\text{where } D_n = \begin{pmatrix} 0 & 0 \\ b_{2n+1} & 0 \end{pmatrix}.$$

Let \mathcal{P} be the vector space of polynomials with complex coefficients. It is well known that, given the recurrence relation (6), there exist two linear moment functionals u^1, u^2 from \mathcal{P} to \mathbb{C} verifying

$$\begin{cases} u^i[z^j P_{2m}] = u^i[z^j P_{2m+1}] = 0, & j = 0, 1, \dots, m-1, i = 1, 2, \\ u^1[z^m P_{2m+1}] = 0 \end{cases} \quad (13)$$

(see [10, Theorem 3.2], see also [5,9]).

We consider the space $\mathcal{P}^2 = \{(q_1, q_2)^T : q_i \text{ polynomial}, i = 1, 2\}$ and the space $\mathcal{M}_{2 \times 2}$ of (2×2) -matrices with complex entries.

Definition 1. If the functionals u^1, u^2 verify (13), then we say that the function $\mathcal{W} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ given by

$$\mathcal{W} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} u^1[q_1] & u^2[q_1] \\ u^1[q_2] & u^2[q_2] \end{pmatrix} \quad (14)$$

is a vector of functionals associated with the recurrence relation (6).

If \mathcal{W} is a vector of functionals associated with the recurrence relation (6), then the following orthogonality relations are verified

$$\mathcal{W}(z^j \mathcal{B}_m) = \mathcal{O}_2, \quad j = 0, 1, \dots, m-1, \quad (15)$$

where \mathcal{O}_2 denotes the 2×2 null matrix.

Definition 2. A function $\mathcal{W} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ verifying (15) is called orthogonality vector of functionals for the recurrence relation (7).

Since the above definitions, any vector of functionals associated with the recurrence relation (6) is always an orthogonality vector of functionals for the recurrence relation (7). As in the scalar case, it is possible to find more than one orthogonality vector of functionals. In fact, given a such function $\mathcal{W} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$, and given any matrix $M \in \mathcal{M}_{2 \times 2}$ it is enough to define \mathcal{W}_M as

$$\mathcal{W}_M \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} := \mathcal{W} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} M \quad (16)$$

for having another orthogonality vector of functionals. In the following, we assume that \mathcal{W} is a fixed vector of functionals associated with the recurrence relation (6) such that $\mathcal{W}(\mathcal{B}_0)$ is an invertible matrix.

We recall that, in (7), the matrix C_0 was arbitrary chosen. In the sequel we assume

$$C_0 = \begin{pmatrix} 1 & 0 \\ -a_1 & 1 \end{pmatrix}.$$

Take $M = (\mathcal{W}(\mathcal{B}_0))^{-1} C_0$ and define

$$\mathcal{U} = \mathcal{W}_M$$

as in (16). Then, $\mathcal{U}(\mathcal{Q}) = \mathcal{W}(\mathcal{Q})(\mathcal{W}(\mathcal{B}_0))^{-1} C_0$ for any $\mathcal{Q} \in \mathcal{P}^2$ and, in particular,

$$\mathcal{U}(\mathcal{B}_0) = C_0. \quad (17)$$

Moreover, from (7) and (15),

$$\mathcal{U}(z^m \mathcal{B}_m) = C_m \mathcal{U}(z^{m-1} \mathcal{B}_{m-1}), \quad m \in \mathbb{N} \quad (18)$$

(see Lemma 1). Using (17), (18), and again (15), for each $m \in \mathbb{N} \cup \{0\}$ we arrive to

$$\mathcal{U}(z^j \mathcal{B}_m) = \begin{cases} \mathcal{O}_2, & j = 0, 1, \dots, m-1, \\ C_m C_{m-1} \cdots C_0, & j = m. \end{cases} \quad (19)$$

We use the vectors $\mathcal{P}_m = \mathcal{P}_m(z) = (z^{2m}, z^{2m+1})^T$ for each $m \in \mathbb{N} \cup \{0\}$. The following definition extends the corresponding to the scalar case.

Definition 3. For each $m = 0, 1, \dots$, the matrix $\mathcal{U}(z^m \mathcal{P}_0)$ is called moment of order m for the vector of functionals \mathcal{U} .

In particular, since $\mathcal{B}_0 = C_0 \mathcal{P}_0$, we have

$$\mathcal{U}(\mathcal{P}_0) = I_2 \quad (20)$$

(see (17)).

We know $\mathcal{U} = \mathcal{U}\{t\}$ depends on t , besides this dependence is not explicitly written (as we said above). Then, it is possible to define the derivative of \mathcal{U} as usual,

$$\frac{d\mathcal{U}}{dt} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$$

such that, for each $\mathcal{B} \in \mathcal{P}^2$,

$$\frac{d\mathcal{U}}{dt}(\mathcal{B}) = \lim_{\Delta t \rightarrow 0} \frac{\mathcal{U}\{t + \Delta t\}(\mathcal{B}) - \mathcal{U}\{t\}(\mathcal{B})}{\Delta t}.$$

Obviously, the usual properties for this kind of operators are verified. In particular,

$$\frac{d}{dt}(\mathcal{U}(\mathcal{B})) = \frac{d\mathcal{U}}{dt}(\mathcal{B}) + \mathcal{U}(\dot{\mathcal{B}}), \quad \forall \mathcal{B} \in \mathcal{P}^2. \quad (21)$$

We use $\frac{d\mathcal{U}}{dt}$ and (21) below. Also, we will use the matrix function \mathcal{R}_J given in (5). We define the *generating function of the moments* as

$$\mathcal{F}_J(z) = C_0^{-1} \mathcal{R}_J(z) C_0, \quad |z| > \|J\|. \quad (22)$$

Next, we have our second main result, related with Theorem 1. More precisely speaking, we will see that Theorem 1 follows directly from Theorem 2.

Theorem 2. In the conditions of Theorem 1, assume $\dot{a}_1 = b_1$. Let \mathcal{U} be given by (19). Then, the following assertions are equivalent:

(e) $\{a_n, b_n, c_n\}, n \in \mathbb{N}$, is a solution of (1), this is, (9) holds.

(f) For each $n = 0, 1, \dots$, we have

$$\frac{d}{dt} \mathcal{U}(z^n \mathcal{P}_0) = \mathcal{U}(z^{n+1} \mathcal{P}_0) - \mathcal{U}(z^n \mathcal{P}_0) \mathcal{U}(z \mathcal{P}_0). \quad (23)$$

(g) For all $\zeta \in \mathbb{C}$ such that $|\zeta| > \|J\|$,

$$\dot{\mathcal{F}}_J(\zeta) = \mathcal{F}_J(\zeta) (\zeta I_2 - \mathcal{U}(z \mathcal{P}_0)) - I_2, \quad (24)$$

being \mathcal{F}_J the generating function defined in (22).

(h) For all $\mathcal{B} \in \mathcal{P}^2$ we have

$$\left(\frac{d}{dt} \mathcal{U} \right) (\mathcal{B}) = \mathcal{U}(z \mathcal{B}) - \mathcal{U}(\mathcal{B}) \mathcal{U}(z \mathcal{P}_0). \quad (25)$$

(i) For each $n = 0, 1, \dots$, we have (12).

Moreover, we have other consequences of Theorem 1 and Theorem 2.

In the next result, $\int_0^t f(s) ds$ is understood to be the solution $X = X(t)$ of the Cauchy problem

$$\left. \begin{array}{l} \dot{X} = f(t), \\ X(0) = 0 \end{array} \right\}$$

in the suitable space. It is well known that, in our conditions, there exists a unique solution of this problem (see, for instance, [6,11]).

Corollary 1. Under the conditions of Theorem 1, if $\{a_n, b_n, c_n\}, n \in \mathbb{N}$, is a solution of (1), then

$$\mathcal{R}_J(z) = \exp(z t) C_0 M(t, z) (N(t))^{-1}, \quad (26)$$

where

$$N(t) = \begin{pmatrix} \exp(\int_0^t a_1 ds) & \exp(\int_0^t a_1 ds) \int_0^t \exp(-\int_0^s (a_2 - a_1) dr) ds \\ 0 & \exp(\int_0^t a_2 ds) \end{pmatrix},$$

$$M(t, z) = - \int_0^t \exp(-zs) C_0^{-1} N(s) ds + (C_0(0))^{-1} \mathcal{R}_0(z)$$

(here, $C_0(0)$ and $\mathcal{R}_0(z)$ are, respectively, C_0 and $\mathcal{R}(z)$ for $t = 0$).

Given a linear functional $u : \mathcal{P} \rightarrow \mathbb{C}$, we may define the new functional $e^{zt}u : \mathcal{P} \rightarrow \mathbb{C}$ as

$$(e^{zt}u)[z^j] = \sum_{k \geq 0} \frac{t^k}{k!} u[z^{k+j}]. \quad (27)$$

We denote by $\mathcal{U}_0 = (u_0^1, u_0^2)^T$ our vector of functionals \mathcal{U} when $t = 0$ and, similarly, by J_0 the triangular matrix given in (2) when $t = 0$. If J_0 is a bounded matrix, then

$$|u_0^i[z^{k+j}]| \leq m_{ij} \|J_0\|^{k+j}, \quad i = 1, 2, k, j = 0, 1, \dots,$$

and the right-hand side of (27) is well defined for $u = u_0^i$, $i = 1, 2$ (see [7, Theorem 4]). In this case, we can define the vector of functionals $e^{zt}\mathcal{U}_0$ as

$$(e^{zt}\mathcal{U}_0)(\mathcal{B}) = \begin{pmatrix} (e^{zt}u_0^1)[q_1] & (e^{zt}u_0^2)[q_1] \\ (e^{zt}u_0^1)[q_2] & (e^{zt}u_0^2)[q_2] \end{pmatrix}$$

for each $\mathcal{B} = (p, q)^T \in \mathcal{P}^2$. As in [3, Theorem 3], we give a possible representation for the vector of functionals \mathcal{U} .

Corollary 2. *In the conditions of Theorem 2, and with the notation used in (16), assume that the vector of functionals \mathcal{U} verifies*

$$\mathcal{U} = (e^{zt}\mathcal{U}_0)_M \quad (28)$$

for some $M \in \mathcal{M}_{2 \times 2}$. Then, $\{a_n, b_n, c_n\}$, $n \in \mathbb{N}$, is a solution of (1).

In Section 2 we show that the study of the system (1) can be reduced to the evolution of the main block of the matrix J^n i.e. J_{11}^n . In Section 3 we prove Theorem 2. The main idea is to express (1) in terms of the evolution of the moments. In Section 4 we prove Theorem 1. The main feature of this result is the connections between the resolvent function, \mathcal{R}_j , and the evolution of the system of vectorial polynomials, $\{\mathcal{B}_n\}$.

2. Auxiliary results

Next lemma was used for proving (18).

Lemma 1. *Let \mathcal{W} be a vector of functionals associated with (6). Then*

$$\mathcal{W}(A_1 \mathcal{Q}_1 + A_2 \mathcal{Q}_2) = A_1 \mathcal{W}(\mathcal{Q}_1) + A_2 \mathcal{W}(\mathcal{Q}_2) \quad (29)$$

is verified for any $\mathcal{Q}_1, \mathcal{Q}_2 \in \mathcal{P}^2$ and $A_1, A_2 \in \mathcal{M}_{2 \times 2}$.

Proof. It is sufficient to take into account that $\mathcal{W} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ is given by

$$\mathcal{W} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \begin{pmatrix} u[q_1] & v[q_1] \\ u[q_2] & v[q_2] \end{pmatrix}$$

when $u, v : \mathcal{P} \rightarrow \mathbb{C}$ are linear functionals. \square

Lemma 2. *The orthogonality vector of functionals $\mathcal{U} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ is determined by (19). This is, \mathcal{U} is the unique vector of functionals associated with the recurrence relation (6) verifying (19).*

Proof. Given $(q_1, q_2)^T \in \mathcal{P}^2$, for each $i = 1, 2$ we can write

$$q_i(z) = \sum_{k=1}^2 \alpha_{ik}^0 P_{k-1}(z) + \sum_{k=1}^2 \alpha_{ik}^1 P_{1+k}(z) + \dots + \sum_{k=1}^2 \alpha_{ik}^m P_{2m+k-1}(z), \quad \alpha_{ik}^j \in \mathbb{C},$$

where $m = \max\{m_1, m_2\}$ and $\deg(q_i) \leq 2m_i + 1$ (we understand $\alpha_{ik}^j = 0$ when $j > m_i$). This is,

$$(q_1, q_2)^T = \sum_{j=0}^m E_j \mathcal{B}_j, \quad (30)$$

being $E_j = (\alpha_{ik}^j) \in \mathcal{M}_{2 \times 2}$, $j = 0, \dots, m$. From (29), if $\tilde{\mathcal{U}} : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ is a vector of functionals associated with the recurrence relation (6), then

$$\tilde{\mathcal{U}} \begin{pmatrix} q_1 \\ q_2 \end{pmatrix} = \sum_{j=0}^m E_j \tilde{\mathcal{U}}(\mathcal{B}_j). \quad (31)$$

Moreover, if $\tilde{\mathcal{U}}$ verifies (19) we have $\tilde{\mathcal{U}}(\mathcal{B}_j) = \mathcal{U}(\mathcal{B}_j) = 0$, $j = 1, 2, \dots$, and also $\tilde{\mathcal{U}}(\mathcal{B}_0) = \mathcal{U}(\mathcal{B}_0) = C_0$. Therefore, from (31) we arrive to $\tilde{\mathcal{U}} = \mathcal{U}$. \square

Next result shows that it is possible to recover the entries of matrix J using the orthogonality vector of functionals \mathcal{U} .

Lemma 3. *The entries of the matrix J are determined by the sequence $\{\mathcal{B}_n\}$ of vectorial polynomials.*

Proof. The entries of J are determined by the blocks C_n, \mathcal{B}_n , $n \in \mathbb{N}$. Then, it is sufficient to express these blocks in terms of $\{\mathcal{B}_n\}$. Since C_k , $k = 0, 1, \dots$, are invertible matrices, also $\mathcal{U}(z^k \mathcal{B}_k)$ is invertible and, from (19),

$$C_n = \mathcal{U}(z^n \mathcal{B}_n) (\mathcal{U}(z^{n-1} \mathcal{B}_{n-1}))^{-1}, \quad n = 1, 2, \dots$$

On the other hand, from (7) and (19) we deduce

$$C_n \mathcal{U}(z^n \mathcal{B}_{n-1}) + B_{n+1} \mathcal{U}(z^n \mathcal{B}_n) - \mathcal{U}(z^{n+1} \mathcal{B}_n) = 0, \quad n = 0, 1, \dots$$

Then, for $n \in \mathbb{N}$ we have

$$B_n = (\mathcal{U}(z^n \mathcal{B}_{n-1}) - C_{n-1} \mathcal{U}(z^{n-1} \mathcal{B}_{n-2})) (\mathcal{U}(z^{n-1} \mathcal{B}_{n-1}))^{-1},$$

and the result follows. \square

Next, we determine the expression of the moment $\mathcal{U}(\mathcal{P}_n) = \mathcal{U}(x^n \mathcal{P}_0)$ in terms of the matrix J .

Lemma 4. *For each $n = 0, 1, \dots$ we have*

$$\mathcal{U}(z^n \mathcal{P}_0) = C_0^{-1} J_{11}^n C_0. \quad (32)$$

Proof. We know that $\mathcal{U}(\mathcal{P}_0) = I_2$ (see (20)), then the result is verified for $n = 0$.

Another way to express (7) is

$$J \begin{pmatrix} \mathcal{B}_0(z) \\ \mathcal{B}_1(z) \\ \vdots \end{pmatrix} = z \begin{pmatrix} \mathcal{B}_0(z) \\ \mathcal{B}_1(z) \\ \vdots \end{pmatrix}.$$

Thus,

$$J^n \begin{pmatrix} \mathcal{B}_0(z) \\ \mathcal{B}_1(z) \\ \vdots \end{pmatrix} = z^n \begin{pmatrix} \mathcal{B}_0(z) \\ \mathcal{B}_1(z) \\ \vdots \end{pmatrix}, \quad n \in \mathbb{N}. \quad (33)$$

Comparing the first rows in (33), and taking into account (4) and the form of J ,

$$\sum_{i \geq 1} J_{1i}^n \mathcal{B}_{i-1}(z) = J_{11}^n \mathcal{B}_0(z) + J_{12}^n \mathcal{B}_1(z) = z^n \mathcal{B}_0(z).$$

Then, from (19),

$$\mathcal{U}(z^n \mathcal{B}_0) = J_{11}^n \mathcal{U}(\mathcal{B}_0),$$

this is,

$$C_0 \mathcal{U}(z^n \mathcal{P}_0) = J_{11}^n C_0$$

(see (17) and (20)), which is (32). \square

The following result concerns to solutions associated with the matrix J , non-necessarily bounded.

Lemma 5. *If the sequence $\{a_n, b_n, c_n\}$, $n \in \mathbb{N}$, is a solution of (1), then (10) is verified.*

Proof. Under the given conditions, we know that (9) holds. Then, it is very easy to verify

$$\frac{d}{dt} J^n = J^n J_- - J_- J^n$$

and, with the established notation,

$$\frac{d}{dt} J_{11}^n = (J^n J_-)_{11} - (J_- J^n)_{11}.$$

From (2) and (4),

$$\left. \begin{aligned} (J^n J_-)_{11} &= J_{11}^n (J_-)_{11} + J_{12}^n (J_-)_{21}, \\ (J_- J^n)_{11} &= (J_-)_{11} J_{11}^n. \end{aligned} \right\}$$

Then,

$$\dot{J}_{11}^n = J_{11}^n (J_-)_{11} - (J_-)_{11} J_{11}^n + J_{12}^n (J_-)_{21}. \quad (34)$$

On the other hand, $J_{11}^{n+1} = J_{11}^n J_{11} + J_{12}^n (J_-)_{21}$, which, joint with (34), goes to (10). \square

3. Proof of Theorem 2

In the first place, we show (e) \Rightarrow (f). Assume that (9) holds. Since Lemma 5, we have (10). Moreover, due to Lemma 4, $J_{11}^n = C_0 \mathcal{U}(z^n \mathcal{P}_0) C_0^{-1}$, $n \in \mathbb{N}$, and, in particular, $B_1 = J_{11} = C_0 \mathcal{U}(z \mathcal{P}_0) C_0^{-1}$. Also, $C_0^{-1} (J_-)_{11} = (J_-)_{11} C_0 = (J_-)_{11}$. Then, from (34),

$$\begin{aligned} \frac{d}{dt} (C_0 \mathcal{U}(z^n \mathcal{P}_0) C_0^{-1}) &= C_0 \mathcal{U}(z^{n+1} \mathcal{P}_0) C_0^{-1} - C_0 \mathcal{U}(z^n \mathcal{P}_0) \mathcal{U}(z \mathcal{P}_0) C_0^{-1} \\ &\quad + C_0 \mathcal{U}(z^n \mathcal{P}_0) (J_-)_{11} - (J_-)_{11} \mathcal{U}(z^n \mathcal{P}_0) C_0^{-1}. \end{aligned} \quad (35)$$

On the other hand, taking derivatives (and denoting by \dot{C}_0^{-1} the derivative of C_0^{-1}),

$$\frac{d}{dt} (C_0 \mathcal{U}(z^n \mathcal{P}_0) C_0^{-1}) = \dot{C}_0 \mathcal{U}(z^n \mathcal{P}_0) C_0^{-1} + C_0 \mathcal{U}(z^n \mathcal{P}_0) \dot{C}_0^{-1} + C_0 \frac{d}{dt} (\mathcal{U}(z^n \mathcal{P}_0)) C_0^{-1}. \quad (36)$$

Since $\dot{a}_1 = b_1$, we can verify $\dot{C}_0 = -(J_-)_{11}$, $\dot{C}_0^{-1} = (J_-)_{11}$. Hence, comparing the right-hand sides of (35) and (36) we arrive to (23) (we recall that C_0 is an invertible matrix).

Now, we prove (f) \Rightarrow (g). Due to (5) and Lemma 4,

$$\mathcal{F}_J(\zeta) = \sum_{n \geq 0} \frac{\mathcal{U}(z^n \mathcal{P}_0)}{\zeta^{n+1}}, \quad |\zeta| > \|J\|. \quad (37)$$

Then, from (23),

$$\begin{aligned} \dot{\mathcal{F}}_J(\zeta) &= \sum_{n \geq 0} \frac{\mathcal{U}(z^{n+1} \mathcal{P}_0) - \mathcal{U}(z^n \mathcal{P}_0) \mathcal{U}(z \mathcal{P}_0)}{\zeta^{n+1}} \\ &= \zeta \sum_{n \geq 0} \frac{\mathcal{U}(z^{n+1} \mathcal{P}_0)}{\zeta^{n+2}} - \sum_{n \geq 0} \frac{\mathcal{U}(z^n \mathcal{P}_0)}{\zeta^{n+1}} \mathcal{U}(z \mathcal{P}_0) \\ &= \zeta \left(\mathcal{F}_J(\zeta) - \frac{1}{\zeta} I_2 \right) - \mathcal{F}_J(\zeta) \mathcal{U}(z \mathcal{P}_0). \end{aligned}$$

This is, (24) is verified.

Given (24), we are going to obtain the derivative of the vector of functionals \mathcal{U} . For doing this, we use the linearity of \mathcal{U} and the convergence of the series,

$$\mathcal{F}_J(\zeta) = \mathcal{U}\left(\sum_{n \geq 0} \frac{z^n}{\zeta^{n+1}} \mathcal{P}_0\right) = \mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right), \quad |\zeta| > \|J\|. \quad (38)$$

(Here and in the next expressions, as usual, $\mathcal{U} = \mathcal{U}_z$ is the vector of functionals defined on the closure of the space \mathcal{P}^2 of vectorial polynomials $(q_1, q_2)^T$ in the variable z .)

From (38) and (24),

$$\begin{aligned} \frac{d}{dt} \mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right) &= \mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right) (\zeta I_2 - \mathcal{U}(z \mathcal{P}_0)) - I_2 \\ &= \mathcal{U}\left(\left(1 + \frac{z}{\zeta - z}\right) \mathcal{P}_0\right) - \mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right) \mathcal{U}(z \mathcal{P}_0) - I_2 \\ &= \mathcal{U}\left(\frac{z}{\zeta - z} \mathcal{P}_0\right) - \mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right) \mathcal{U}(z \mathcal{P}_0). \end{aligned} \quad (39)$$

Define the vectors of functionals $\mathcal{U}_1, \mathcal{U}_2 : \mathcal{P}^2 \rightarrow \mathcal{M}_{2 \times 2}$ such that

$$\left. \begin{aligned} \mathcal{U}_1(\mathcal{B}) &= \mathcal{U}(z \mathcal{B}), \\ \mathcal{U}_2(\mathcal{B}) &= \mathcal{U}(\mathcal{B}) \mathcal{U}(z \mathcal{P}_0) \end{aligned} \right\} \quad (40)$$

for each $\mathcal{B} \in \mathcal{P}^2$. We remark that $\frac{1}{\zeta - z} \mathcal{P}_0$ do not depend on $t \in \mathbb{R}$. In (39), denoting $\dot{\mathcal{U}} = \frac{d}{dt} \mathcal{U}$, we have $\dot{\mathcal{U}} = \mathcal{U}_1 - \mathcal{U}_2$ over $\frac{1}{\zeta - z} \mathcal{P}_0$, being

$$\mathcal{U}\left(\frac{1}{\zeta - z} \mathcal{P}_0\right) = \frac{1}{\zeta} \mathcal{U}(\mathcal{P}_0) + \frac{1}{\zeta^2} \mathcal{U}(z \mathcal{P}_0) + \dots, \quad |\zeta| > \|J\|.$$

Hence, we have $\dot{\mathcal{U}} = \mathcal{U}_1 - \mathcal{U}_2$ over \mathcal{P}^2 , this is, we have (25).

For proving (h) \Rightarrow (i), as in (30), $\dot{\mathcal{B}}_n$ can be written in terms of the sequence $\{\mathcal{B}_n\}$,

$$\dot{\mathcal{B}}_n(z) = D_0^{(n)} \mathcal{B}_0(z) + D_1^{(n)} \mathcal{B}_1(z) + \dots + D_n^{(n)} \mathcal{B}_n(z). \quad (41)$$

If $n = 0, 1$, the above expression is

$$\dot{\mathcal{B}}_n(z) = D_{n-1}^{(n)} \mathcal{B}_{n-1}(z) + D_n^{(n)} \mathcal{B}_n(z). \quad (42)$$

Let $n \geq 2$ be fixed. We are going to show that (42) holds, also, for n . Due to the orthogonality, from (41),

$$\mathcal{U}(\dot{\mathcal{B}}_n) = D_0^{(n)} \mathcal{U}(\mathcal{B}_0).$$

In fact, using (25),

$$\begin{aligned} \mathcal{O}_2 &= \frac{d}{dt} (\mathcal{U}(\mathcal{B}_n)) = \dot{\mathcal{U}}(\mathcal{B}_n) + \mathcal{U}(\dot{\mathcal{B}}_n) \\ &= \mathcal{U}(z \mathcal{B}_n) - \mathcal{U}(\mathcal{B}_n) \mathcal{U}(z \mathcal{P}_0) + D_0^{(n)} \mathcal{U}(\mathcal{B}_0). \end{aligned}$$

Thus, $D_0^{(n)} = \mathcal{O}_2$. We proceed by induction on n , assuming

$$D_0^{(n)} = \dots = D_{j-1}^{(n)} = \mathcal{O}_2$$

for a fixed $j < n - 1$. Using (41) and, again, (25) and (19),

$$\begin{aligned} \mathcal{O}_2 &= \frac{d}{dt} \mathcal{U}(z^j \mathcal{B}_n) = \left(\frac{d}{dt} \mathcal{U}\right)(z^j \mathcal{B}_n) + \mathcal{U}(z^j \dot{\mathcal{B}}_n) \\ &= \mathcal{U}(z^{j+1} \mathcal{B}_n) - \mathcal{U}(z^j \mathcal{B}_n) \mathcal{U}(z \mathcal{P}_0) + D_j^{(n)} \mathcal{U}(z^j \mathcal{B}_j) \\ &= D_j^{(n)} \mathcal{U}(z^j \mathcal{B}_j), \end{aligned}$$

where $\mathcal{U}(z^j \mathcal{B}_j)$ is an invertible matrix. Thus, $D_j^{(n)} = \mathcal{O}_2$ and (42) is verified for any $n \in \mathbb{N}$.

Our next purpose is to determine $D_j^{(n)}$, $j = n - 1, n$. From (42),

$$\mathcal{U}(z^{n-1}\dot{\mathcal{B}}_n) = D_{n-1}^{(n)}\mathcal{U}(z^{n-1}\mathcal{B}_{n-1}).$$

Then, because of (25) and (19),

$$\mathcal{O}_2 = \frac{d}{dt}\mathcal{U}(z^{n-1}\mathcal{B}_n) = \mathcal{U}(z^n\mathcal{B}_n) - \mathcal{U}(z^{n-1}\mathcal{B}_n)\mathcal{U}(z\mathcal{P}_0) + D_{n-1}^{(n)}\mathcal{U}(z^{n-1}\mathcal{B}_{n-1})$$

and, therefore,

$$D_{n-1}^{(n)} = -C_n.$$

On the other hand, writing

$$\mathcal{B}_n(z) = \sum_{i=0}^n F_i^{(n)}\mathcal{P}_i(z) \quad (43)$$

and comparing the coefficient of z^{2n} and z^{2n+1} in both sides of (43), we obtain

$$F_n^{(n)} = \begin{pmatrix} 1 & 0 \\ f_n & 1 \end{pmatrix}.$$

Moreover, taking derivatives in (43) and comparing with (42), we see $D_n^{(n)} = \dot{F}_n^{(n)}$ or, what is the same,

$$D_n^{(n)} = \begin{pmatrix} 0 & 0 \\ d_n & 0 \end{pmatrix},$$

where we need to determine d_n . From (25) and (42),

$$D_n^{(n)}\mathcal{U}(z^n\mathcal{B}_n) = \left(\frac{d}{dt}\mathcal{U}(z^n\mathcal{B}_n) - \mathcal{U}(z^{n+1}\mathcal{B}_n) + \mathcal{U}(z^n\mathcal{B}_n)\mathcal{U}(z\mathcal{P}_0) \right) + C_n\mathcal{U}(z^n\mathcal{B}_{n-1}).$$

Then, using (19) and (7),

$$D_n^{(n)} = \left(\frac{d}{dt}(C_n C_{n-1} \cdots C_0) \right) (C_n C_{n-1} \cdots C_0)^{-1} - B_{n+1} + (C_n C_{n-1} \cdots C_0)\mathcal{U}(z\mathcal{P}_0)(C_n C_{n-1} \cdots C_0)^{-1},$$

thus

$$D_n^{(n)} + B_{n+1} = \left(\frac{d}{dt}(C_n C_{n-1} \cdots C_1) \right) (C_n C_{n-1} \cdots C_1)^{-1} + (C_n C_{n-1} \cdots C_1)(\dot{C}_0 C_0^{-1} + J_{11})(C_n C_{n-1} \cdots C_1)^{-1}. \quad (44)$$

The matrix $C_n C_{n-1} \cdots C_1$ is upper triangular. Moreover, because of $\dot{a}_1 = b_1$ also $\dot{C}_0 C_0^{-1} + J_{11}$ is an upper triangular matrix and, then, the matrix in the left-hand side of (44) is upper triangular and, consequently, $d_n = b_{2n+1}$.

Finally, we show (i) \Rightarrow (e). Taking derivatives in (7),

$$\dot{C}_n \mathcal{B}_{n-1}(z) + \dot{B}_{n+1} \mathcal{B}_n(z) + C_n \dot{\mathcal{B}}_{n-1}(z) + (B_{n+1} - zI_2) \dot{\mathcal{B}}_n(z) + A \dot{\mathcal{B}}_{n+1}(z) = 0, \quad n = 0, 1, \dots$$

Using (12) and taking into account $AD_{n+1} = D_n A = \mathcal{O}_2$,

$$\begin{aligned} & \dot{C}_n \mathcal{B}_{n-1}(z) + \dot{B}_{n+1} \mathcal{B}_n(z) + C_n (-C_{n-1} \mathcal{B}_{n-2}(z) + D_{n-1} \mathcal{B}_{n-1}(z)) \\ & + (B_{n+1} - zI_2) (-C_n \mathcal{B}_{n-1}(z) + D_n \mathcal{B}_n(z)) - AC_{n+1} \mathcal{B}_n(z) = 0. \end{aligned} \quad (45)$$

Using, again, (7) for eliminating the explicit expression in z ,

$$\left. \begin{aligned} z\mathcal{B}_n(z) &= C_n \mathcal{B}_{n-1}(z) + B_{n+1} \mathcal{B}_n(z) + A \mathcal{B}_{n+1}(z), \\ z\mathcal{B}_{n-1}(z) &= C_{n-1} \mathcal{B}_{n-2}(z) + B_n \mathcal{B}_{n-1}(z) + A \mathcal{B}_n(z). \end{aligned} \right\}$$

Substituting in (45), and identifying with zero the coefficients of the vectorial polynomials in the obtained expression, we arrive to

$$\left. \begin{aligned} \dot{B}_n &= AC_n - C_{n-1}A + D_{n-1}B_n - B_n D_{n-1}, \\ \dot{C}_n &= D_n C_n - C_n D_{n-1} + B_{n+1}C_n - C_n B_n, \end{aligned} \right\} \quad n = 1, 2, \dots \quad (46)$$

Taking into account that, with the above notation, $D_n = (J_-)_{n+1, n+1}$, we see that (46) is equivalent to (9) when we consider J as a blocked matrix.

4. Proof of Theorem 1 and corollaries

4.1. Proof of Theorem 1

We start by establishing the equivalence between (10) and (11). The key is the convergence in the respective operator norm, for $|z| > \|J\|$, of the series given in the right-hand side of (3) and (5). Starting by (10), to obtain $\dot{\mathcal{R}}_J(z)$ it is sufficient to take derivatives in (5) and to substitute j_{11}^n in

$$\dot{\mathcal{R}}_J(z) = \sum_{n \geq 0} \frac{j_{11}^n}{z^{n+1}}, \quad |z| > \|J\|.$$

Reciprocally, if (11) holds, substituting $\mathcal{R}_J(z)$ and $\dot{\mathcal{R}}_J(z)$ by its Laurent expansion, and comparing their coefficients, we arrive to (10).

The rest of the proof is to show the equivalence between (a), (b) and (d).

(a) \Rightarrow (b) is Lemma 5.

Now, we are going to prove (b) \Rightarrow (d). We assume that (10) is verified. Taking $n = 1$ in this expression we immediately deduce $\dot{a}_1 = b_1$. Moreover, from (10) we arrive to (23) in the same way that in the proof of (e) \Rightarrow (f) in Theorem 2. Then we are under the hypothesis of Theorem 2 and, therefore, we have (12).

Finally, (12) \Rightarrow (9) was proved in Theorem 2.

4.2. Proof of Corollary 1

It is easy to see that C_0 , $M(t, z)$ and $N(t)$ are, respectively, the solutions of the following Cauchy problems:

$$\left. \begin{aligned} \dot{X} &= -(J_-)_{11}X, \\ X(0) &= C_0(0), \end{aligned} \right\} \quad \left. \begin{aligned} \dot{X} &= -\exp(-zt)C_0^{-1}N(t), \\ X(0) &= (C_0(0))^{-1}\mathcal{R}_0(z), \end{aligned} \right\}$$

and

$$\left. \begin{aligned} \dot{X} &= (J_{11} - (J_-)_{11})X, \\ X(0) &= I_2. \end{aligned} \right\}$$

Taking derivatives in the right-hand side of (26), and checking the initial condition, we can prove that $\exp(zt)C_0M(t, z) \times (N(t))^{-1}$ is a solution of the following Cauchy problem,

$$\left. \begin{aligned} \dot{X} &= X(zI_2 - J_{11}) - I_2 + [X, (J_-)_{11}], \\ X(0) &= \mathcal{R}_0(z). \end{aligned} \right\} \quad (47)$$

From [11], we know that (47) has a unique solution. On the other hand, since Theorem 1, $\mathcal{R}(z)$ is solution of (47). Then, we arrive to (26).

4.3. Proof of Corollary 2

Since (20), in the conditions of Corollary 2, necessarily (28) implies

$$M = [(e^{zt}\mathcal{U}_0)(\mathcal{P}_0)]^{-1}.$$

On the other hand, for proving that $\{a_n, b_n, c_n\}$, $n \in \mathbb{N}$, is a solution of (1), it is sufficient to show that (23) holds. Let $k \in \{0, 1, \dots\}$ be fixed. Taking into account

$$\frac{d}{dt}(e^{zt}\mathcal{U}_0)(z^k\mathcal{P}_0) = (e^{zt}\mathcal{U}_0)(z^{k+1}\mathcal{P}_0)$$

and

$$\frac{d}{dt}M = -MU(z\mathcal{P}_0),$$

and taking derivatives in

$$\mathcal{U}(z^k\mathcal{P}_0) = (e^{zt}\mathcal{U}_0)(z^k\mathcal{P}_0)M$$

(see (28)), we arrive to (23).

Acknowledgments

This research was carried out while on a visit of the first author at University of Coimbra. The research of this author was supported in part by research grants MTM2009-14668-C02-02 and PT2009-0031 of "Ministerio de Ciencia e Innovación", Spain. The work of the second author was supported by CMUC/FCT. The third author would like to thank CIDMA–Center for Research and Development in Mathematics and Applications. The second and third authors were supported by "Acção Integrada Luso-Espanhola" n. E-4/10.

References

- [1] A. Aptekarev, V. Kaliaguine, J. Van Iseghem, The genetic sums' representation for the moments of a system of Stieltjes functions and its application, *Constr. Approx.* 16 (4) (2000) 487–524.
- [2] D. Barrios Rolanía, A. Branquinho, Complex high order Toda lattices, *J. Difference Equ. Appl.* 15 (2) (2009) 197–213.
- [3] D. Barrios Rolanía, A. Branquinho, A. Foulquié Moreno, Dynamics and interpretation of some integrable systems via multiple orthogonal polynomials, *J. Math. Anal. Appl.* 361 (2) (2010) 358–370.
- [4] N.M. Ercolani, K.T.-R. McLaughlin, Asymptotics and integrable structures for biorthogonal polynomials associated to a random two-matrix model, in: *Advances in Nonlinear Mathematics and Science*, Phys. D 152/153 (2001) 232–268.
- [5] M.E.H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia Math. Appl., vol. 98, Cambridge Univ. Press, Cambridge, 2005.
- [6] V.G. Jank, Matrix Riccati differential equations, in: A.P. Santana, J.S. Neves, M.P. Oliveira (Eds.), *Textos Mat.*, vol. 36, DMUC, Coimbra, 2005.
- [7] V. Kaliaguine, The operator moment problem, vector continued fractions and an explicit form of the Favard theorem for vector orthogonal polynomials, *J. Comput. Appl. Math.* 65 (1–3) (1995) 181–193.
- [8] V. Sorokin, J. Van Iseghem, Matrix Hermite–Padé problem and dynamical systems, *J. Comput. Appl. Math.* 122 (1–2) (2000) 275–295.
- [9] W. Van Assche, Nonsymmetric linear difference equations for multiple orthogonal polynomials, in: D. Levi, O. Ragnisco (Eds.), *SIDE II – Symmetries and Integrability of Difference Equations*, in: CRM Proc. Lecture Notes, vol. 25, Amer. Math. Soc., 2000, pp. 415–429.
- [10] J. Van Iseghem, Vector orthogonal relations. Vector QD-algorithm, *J. Comput. Appl. Math.* 19 (1) (1987) 141–150.
- [11] N.Ya. Vilenkin, et al., *Functional Analysis*, Wolters-Noordhoff, The Netherlands, 1972.
- [12] K. Yosida, *Functional Analysis*, Springer-Verlag, Berlin, 1995.